Connexions module: m10419

## VECTOR SPACE\*

## Phil Schniter

This work is produced by The Connexions Project and licensed under the Creative Commons Attribution License  $^\dagger$ 

## Abstract

This module introduces vector space.

- A vector space consists of the following four elements:
  - a. A set of vectors V,
  - b. A field of scalars  $\mathbb{F}$  (where, for our purposes,  $\mathbb{F}$  is either  $\mathbb{R}$  or  $\mathbb{C}$ ),
  - c. The operations of vector addition "+" (i.e.,  $+: V \times V \rightarrow V$ )
  - d. The operation of scalar multiplication "."  $(i.e., \cdot : \mathbb{F} \times V \to V)$

for which the following properties hold. (Assume  $x \wedge y \wedge z \in V$  and  $\alpha \wedge \beta \in \mathbb{F}$ .)

Properties	Examples
commutativity	x + y = y + x
associativity –	(x+y) + z = x + (y+z)
	$(\alpha\beta)x = \alpha(\beta x)$
distributivity –	$\alpha \cdot (x+y) = (\alpha \cdot x) + (\alpha \cdot y)$
	$(\alpha + \beta) x = \alpha x + \beta x$
additive identity	$\forall x \in V : (\exists 0, 0 \in V : (x + 0 = x))$
additive inverse	$\forall x \in V : (\exists -x, (-x) \in V : (x + -x = 0))$
multiplicative identity	$\forall x \in V : (1 \cdot x = x)$

Table 1

Important examples of vector spaces include

Properties	Examples
real $N$ -vectors	$V=\mathbb{R}^N,\mathbb{F}=\mathbb{R}$
complex $N$ -vectors	$V=\mathbb{C}^N,\mathbb{F}=\mathbb{C}$
sequences in " $l_p$ "	$V = \left\{ x [n] \mid \exists n \in \mathbb{Z} : \left( \sum_{n=-\infty}^{\infty} ( x[n] )^p < \infty \right) \right\}, \mathbb{F} = \mathbb{C}$
functions in " $\mathcal{L}_p$ "	$V = \left\{ f(t) \mid \int_{-\infty}^{\infty} ( f(t) )^{p} dt < \infty \right\},  \mathbb{F} = \mathbb{C}$

<sup>\*</sup>Version 2.13: Oct 18, 2005 4:34 pm GMT-5

 $<sup>^\</sup>dagger {\rm http://creative commons.org/licenses/by/1.0}$ 

Connexions module: m10419 2

## Table 2

where we have assumed the usual definitions of addition and multiplication. From now on, we will denote the arbitrary vector space  $(V, \mathbb{F}, +, \cdot)$  by the shorthand V and assume the usual selection of  $(\mathbb{F}, +, \cdot)$ . We will also suppress the "·" in scalar multiplication, so that  $\alpha \cdot x$  becomes  $\alpha x$ .

• A subspace of V is a subset  $M \subset V$  for which

a. 
$$\forall x, y, x \in M \land y \in M : ((x + y) \in M)$$
  
b.  $\forall x \in M \land \alpha \in \mathbb{F} : (\alpha x \in M)$ 

NOTE: Note that every subspace must contain 0, and that V is a subspace of itself.

• The **span** of set  $S \subset V$  is the subspace of V containing all linear combinations of vectors in S. When  $S = \{x_0, \ldots, x_{N-1}\},$ 

$$\operatorname{span}(S) := \left\{ \sum_{i=0}^{N-1} \alpha_i x_i \mid \alpha_i \in \mathbb{F} \right\}$$

- A subset of linearly-independent vectors  $\{x_0, \ldots, x_{N-1}\} \subset V$  is called a **basis for V** when its span equals V. In such a case, we say that V has **dimension**N. We say that V is **infinite-dimensional**<sup>1</sup> if it contains an infinite number of linearly independent vectors.
- V is a **direct sum** of two subspaces M and N, written  $V = M \oplus N$ , iff every  $x \in V$  has a **unique** representation x = m + n for  $m \in M$  and  $n \in N$ .

NOTE: Note that this requires  $M \cap N = \{0\}$ 

 $<sup>^{1}</sup>$ The definition of an infinite-dimensional basis would be complicated by issues relating to the convergence of infinite series. Hence we postpone discussion of infinite-dimensional bases until the Hilbert Space (<http://cnx.org/content/m10434/latest/>) section.