Connexions module: m16999

Hypothesis Testing of Single Mean and Single Proportion: Additional Information*

Susan Dean Barbara Illowsky, Ph.D.

This work is produced by The Connexions Project and licensed under the Creative Commons Attribution License †

- In a **hypothesis test** problem, you may see words such as "the level of significance is 1%." The "1%" is the preconceived or preset α .
- The statistician setting up the hypothesis test selects the value of α to use **before** collecting the sample data.
- If no level of significance is given, the accepted standard is to use $\alpha = 0.05$.
- When you calculate the **p-value** and draw the picture, the p-value is the area in the left tail, the right tail, or split evenly between the two tails. For this reason, we call the hypothesis test left, right, or two tailed.
- The alternate hypothesis, H_a , tells you if the test is left, right, or two-tailed. It is the key to conducting the appropriate test.
- H_a **never** has a symbol that contains an equal sign.
- Thinking about the meaning of thep-value: A data analyst (and anyone else) should have more confidence that he made the correct decision to reject the null hypothesis with a smaller p-value (for example, 0.001 as opposed to 0.04) even if using the 0.05 level for alpha. Similarly, for a large p-value like 0.4, as opposed to a p-value of 0.056 (alpha = 0.05 is less than either number), a data analyst should have more confidence that she made the correct decision in failing to reject the null hypothesis. This makes the data analyst use judgment rather than mindlessly applying rules.

The following examples illustrate a left, right, and two-tailed test.

Example 1


 H_o $\mu = 5$ H_a $\mu < 5$

Test of a single population mean. H_a tells you the test is left-tailed. The picture of the p-value is as follows:

^{*}Version 1.13: Jun 13, 2012 5:34 am -0500

[†]http://creativecommons.org/licenses/by/3.0/


Connexions module: m16999 2



Example 2

 $H_o: p \le 0.2$ $H_a: p > 0.2$


This is a test of a single population proportion. H_a tells you the test is **right-tailed**. The picture of the p-value is as follows:



Example 3

 H_o : μ = 50 H_a : μ \neq 50

This is a test of a single population mean. H_a tells you the test is **two-tailed**. The picture of the p-value is as follows.



Glossary

Definition 1: Hypothesis Testing

Based on sample evidence, a procedure to determine whether the hypothesis stated is a reasonable statement and cannot be rejected, or is unreasonable and should be rejected.

Connexions module: m16999 3

Definition 2: p-value

The probability that an event will happen purely by chance assuming the null hypothesis is true. The smaller the p-value, the stronger the evidence is against the null hypothesis.