ENERGY IN WAVES: INTENSITY^{*}

OpenStax College

This work is produced by OpenStax-CNX and licensed under the Creative Commons Attribution License 3.0^{\dagger}

Abstract

• Calculate the intensity and the power of rays and waves.



Figure 1: The destructive effect of an earthquake is palpable evidence of the energy carried in these waves. The Richter scale rating of earthquakes is related to both their amplitude and the energy they carry. (credit: Petty Officer 2nd Class Candice Villarreal, U.S. Navy)

All waves carry energy. The energy of some waves can be directly observed. Earthquakes can shake whole cities to the ground, performing the work of thousands of wrecking balls.

Loud sounds pulverize nerve cells in the inner ear, causing permanent hearing loss. Ultrasound is used for deep-heat treatment of muscle strains. A laser beam can burn away a malignancy. Water waves chew up beaches.

^{*}Version 1.6: Jul 5, 2012 9:06 pm -0500

[†]http://creativecommons.org/licenses/by/3.0/

2

The amount of energy in a wave is related to its amplitude. Large-amplitude earthquakes produce large ground displacements. Loud sounds have higher pressure amplitudes and come from larger-amplitude source vibrations than soft sounds. Large ocean breakers churn up the shore more than small ones. More quantitatively, a wave is a displacement that is resisted by a restoring force. The larger the displacement x, the larger the force F = kx needed to create it. Because work W is related to force multiplied by distance (Fx) and energy is put into the wave by the work done to create it, the energy in a wave is related to amplitude. In fact, a wave's energy is directly proportional to its amplitude squared because

$$W \propto F \mathbf{x} = \mathbf{k} \mathbf{x}^2. \tag{1}$$

The energy effects of a wave depend on time as well as amplitude. For example, the longer deep-heat ultrasound is applied, the more energy it transfers. Waves can also be concentrated or spread out. Sunlight, for example, can be focused to burn wood. Earthquakes spread out, so they do less damage the farther they get from the source. In both cases, changing the area the waves cover has important effects. All these pertinent factors are included in the definition of **intensity** *I* as power per unit area:

$$I = \frac{P}{A} \tag{2}$$

where P is the power carried by the wave through area A. The definition of intensity is valid for any energy in transit, including that carried by waves. The SI unit for intensity is watts per square meter (W/m^2) . For example, infrared and visible energy from the Sun impinge on Earth at an intensity of $1300 W/m^2$ just above the atmosphere. There are other intensity-related units in use, too. The most common is the decibel. For example, a 90 decibel sound level corresponds to an intensity of $10^{-3} W/m^2$. (This quantity is not much power per unit area considering that 90 decibels is a relatively high sound level. Decibels will be discussed in some detail in a later chapter.

Example 1: Calculating intensity and power: How much energy is in a ray of sunlight?

The average intensity of sunlight on Earth's surface is about 700 W/m^2 .

(a) Calculate the amount of energy that falls on a solar collector having an area of 0.500 m^2 in 4.00 h.

(b) What intensity would such sunlight have if concentrated by a magnifying glass onto an area 200 times smaller than its own?

Strategy a

Because power is energy per unit time or $P = \frac{E}{t}$, the definition of intensity can be written as $I = \frac{P}{A} = \frac{E/t}{A}$, and this equation can be solved for E with the given information. Solution a

t:

1. Begin with the equation that states the definition of intensity:

$$I = \frac{P}{A}.$$
(3)

2. Replace P with its equivalent E/

$$I = \frac{E/t}{A}.$$
(4)

3. Solve for E:

$$E = IAt.$$
(5)

4. Substitute known values into the equation:

$$E = \left(700 \text{ W/m}^2\right) \left(0.500 \text{ m}^2\right) \left[(4.00 \text{ h}) (3600 \text{ s/h})\right].$$
(6)

5. Calculate to find E and convert units:

$$5.04 \times 10^6 \text{ J},$$
 (7)

Discussion a

The energy falling on the solar collector in 4 h in part is enough to be useful—for example, for heating a significant amount of water.

Strategy b

Taking a ratio of new intensity to old intensity and using primes for the new quantities, we will find that it depends on the ratio of the areas. All other quantities will cancel.

Solution b

1. Take the ratio of intensities, which yields:

$$\frac{I\prime}{I} = \frac{P\prime/A\prime}{P/A} = \frac{A}{A\prime}$$
 (The powers cancel because $P\prime = P$). (8)

2. Identify the knowns:

$$A = 200A\prime,\tag{9}$$

$$\frac{I\prime}{I} = 200. \tag{10}$$

3. Substitute known quantities:

$$I' = 200I = 200 \left(700 \text{ W/m}^2\right).$$
(11)

4. Calculate to find I':

$$II = 1.40 \times 10^{-5} \,\mathrm{W/m^{-2}}.$$
 (12)

Discussion b

Decreasing the area increases the intensity considerably. The intensity of the concentrated sunlight could even start a fire.

Example 2: Determine the combined intensity of two waves: Perfect constructive interference

If two identical waves, each having an intensity of 1.00 W/m^2 , interfere perfectly constructively, what is the intensity of the resulting wave?

Strategy

We know from Superposition and Interference¹ that when two identical waves, which have equal amplitudes X, interfere perfectly constructively, the resulting wave has an amplitude of 2X. Because a wave's intensity is proportional to amplitude squared, the intensity of the resulting wave is four times as great as in the individual waves.

Ι

Solution

- 1. Recall that intensity is proportional to amplitude squared.
- 2. Calculate the new amplitude:

$$I' \propto (X')^2 = (2X)^2 = 4X^2.$$
(13)

3. Recall that the intensity of the old amplitude was:

$$\propto X^2$$
. (14)

¹"Superposition and Interference" http://cnx.org/content/m42249/latest/

4. Take the ratio of new intensity to the old intensity. This gives:

$$\frac{I\prime}{I} = 4. \tag{15}$$

5. Calculate to find I':

$$I' = 4I = 4.00 \text{ W/m}^2.$$
(16)

Discussion

The intensity goes up by a factor of 4 when the amplitude doubles. This answer is a little disquieting. The two individual waves each have intensities of 1.00 W/m^2 , yet their sum has an intensity of 4.00 W/m^2 , which may appear to violate conservation of energy. This violation, of course, cannot happen. What does happen is intriguing. The area over which the intensity is 4.00 W/m^2 is much less than the area covered by the two waves before they interfered. There are other areas where the intensity is zero. The addition of waves is not as simple as our first look in Superposition and Interference² suggested. We actually get a pattern of both constructive interference and destructive interference whenever two waves are added. For example, if we have two stereo speakers putting out 1.00 W/m^2 each, there will be places in the room where the intensity is 4.00 W/m^2 , other places where the intensity is zero, and others in between. Figure 2 shows what this interference might look like. We will pursue interference patterns elsewhere in this text.



Figure 2: These stereo speakers produce both constructive interference and destructive interference in the room, a property common to the superposition of all types of waves. The shading is proportional to intensity.

1: Check Your Understanding

Which measurement of a wave is most important when determining the wave's intensity?

Solution

Amplitude, because a wave's energy is directly proportional to its amplitude squared.

²"Superposition and Interference" < http://cnx.org/content/m42249/latest/>

1 Section Summary

Intensity is defined to be the power per unit area: $I = \frac{P}{4}$ and has units of W/m².

2 Conceptual Questions

Exercise 2

Two identical waves undergo pure constructive interference. Is the resultant intensity twice that of the individual waves? Explain your answer.

Exercise 3

Circular water waves decrease in amplitude as they move away from where a rock is dropped. Explain why.

3 Problems & Exercises

Exercise 4

Medical Application

Ultrasound of intensity $1.50 \times 10^2 \text{ W/m}^2$ is produced by the rectangular head of a medical imaging device measuring 3.00 by 5.00 cm. What is its power output?

Exercise 5

The low-frequency speaker of a stereo set has a surface area of 0. 05 m² and produces 1W of acoustical power. What is the intensity at the speaker? If the speaker projects sound uniformly in all directions, at what distance from the speaker is the intensity 0. 1 W/m²?

Exercise 6

To increase intensity of a wave by a factor of 50, by what factor should the amplitude be increased?

Exercise 7

Engineering Application

A device called an insolation meter is used to measure the intensity of sunlight has an area of 100 cm^2 and registers 6.50 W. What is the intensity in W/m⁻²?

Exercise 8

Astronomy Application

Energy from the Sun arrives at the top of the Earth's atmosphere with an intensity of 1.30 kW/m^{-2} . How long does it take for $1.8 \times 10^9 \text{ J}$ to arrive on an area of 1. 00 m⁻²?

Exercise 9

Suppose you have a device that extracts energy from ocean breakers in direct proportion to their intensity. If the device produces 10.0 kW of power on a day when the breakers are 1.20 m high, how much will it produce when they are 0.600 m high?

Exercise 10

Engineering Application

(a) A photovoltaic array of (solar cells) is 10.0% efficient in gathering solar energy and converting it to electricity. If the average intensity of sunlight on one day is 700 W/m^{-2} , what area should your array have to gather energy at the rate of 100 W? (b) What is the maximum cost of the array if it must pay for itself in two years of operation averaging 10.0 hours per day? Assume that it earns money at the rate of 9.00 c per kilowatt-hour.

(Solution on p. 7.)

(Solution on p. 7.)

(Solution on p. 7.)

(Solution on p. 7.)

Exercise 11

(Solution on p. 7.)

A microphone receiving a pure sound tone feeds an oscilloscope, producing a wave on its screen. If the sound intensity is originally 2.00×10^{-5} W/m⁻², but is turned up until the amplitude increases by 30.0%, what is the new intensity?

Exercise 12

Medical Application

(a) What is the intensity in W/m² of a laser beam used to burn away cancerous tissue that, when 90.0% absorbed, puts 500 J of energy into a circular spot 2.00 mm in diameter in 4.00 s? (b) Discuss how this intensity compares to the average intensity of sunlight (about 1 W/m^2) and the implications that would have if the laser beam entered your eye. Note how your answer depends on the time duration of the exposure.

Solutions to Exercises in this Module

Solution to Exercise (p. 5) 0.225 WSolution to Exercise (p. 5) 7.07Solution to Exercise (p. 5) 16.0 dSolution to Exercise (p. 5) 2.50 kWSolution to Exercise (p. 6) $3.38 \times 10^{-5} \text{ W/m}^2$

Glossary

Definition 1: intensity power per unit area